

KIR measurement area



Kiruna magnetometers

(1) Vector field (every 1 sec): Fluxgate variometer (Kiruna)

Primary



Secondary

- (3) Total field (every 3-4 sec): Proton magnetometer = scalar value (Kiruna)
- # Primary + Secondary # To be upgraded to 1-sec resolution

#1 spare



(4) Higher frequency than 1 Hz: Pulsation magnetometer (Kiruna)

3 sensors (XYZ) in operation

(2) Absolute measurement (weekly): DI-flux magnetometer (Kiruna)

in operation # spare





Specialty of Kiruna:

- 1) near the iron mining ($\Delta D \sim 10^{\circ}$)
- 2) can combine with other ionospheric measurements
 - auroral image
 - riometer
 - ionosonde
 - mm wave for O₂+O₃ (Zeeman effect)



magnetic observatory facilities (Kiruna)






Highway (E10) effect

Observed effect ~ only 1 nT (at 80 m distance, 90 km/h)

⇒ we do not remove such small spikes



5



Hässleholm (TOP) instrument

Variometer close to Tormestorp

https://www.spaceweather.se/forecast/mag

https://www2.irf.se/mag/

- Site test with DTU: 2013
- Installation and first measurements: 2016
- Real-time since 2018
- Solar observations with DTU 20230502
- Broken spring 2023 => need for backup
- Having backup under discussion



Tormestorp (56°N)

Kiruna (68°N)

Kiruna Suspended Magnetometer Tormestorp magnetometer 2023-08-19 00:00:00 - 2023-08-19 23:59:59 [UTC] 2023-08-19 00:00:00 - 2023-08-19 23:59:59 [UTC] 16900.0 10650 350 52450 16800.0 10600 300 52400 10550 52350 16700.0 10500 52300 **52250** ⊨ 16600.0 10450 **□** 10400 52200 16500.0 10350 52150 10300 52100 16400.0 10250 52050 10200 16300.0 52000 10150 51950 00 01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23 00 01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23



1600.0 r 48200.0

1500.0 - 48100.0

1400.0 - 48000.0

1300.0 - 47900.0 -

1200.0 - 47800.0

1100.0 - 47700.0

1000.0 L 47600.0


Recent developments

- Keep improving our data quality (hardware)
 - 2020: UPS house, true north, replace main variometer
 - 2023: New 1 Hz portable scalar magnetometer is coming
- Keep improving our data quality (software)
 - 2022: Start estimating magpy, scale factor (baseline Z)
 - 2023: Improved set of data 2021, 2022 are ready
- Local |B| anomaly near KIR station
 - Retrieved past past measurements by SGU
 - To be further examined (with new magnetometer)
- Thunderstorm (2023-6-29: strongest? since 2014)
 - Damage on primary proton and data correcting system
 - Damage was within level of quick fix (same as 2022-7-1)
- Possible fear: E10 highway broadening (work starts 2027~)




|B| anomaly ~200 m resolution (by SGU)

- KIR site sits at nearly one extreme (-440 nT).
- Deviation is nearly 3000 nT (out of Z ~ 52000 nT & H ~ 10000 nT)
- Old measurements with 30 km reslution show ΔD ~ 8° around KIR
- We plan more local measurement with portable scalar magnetometer coming soon




Lightening 2023-6-29 (recover 2023-7-03)



E10 will be widened ~ double (2+1 lane)

work to be started 2027



Recent developments (summary)


- Keep improving our data quality (hardware + software)
- |B| anomaly and D anomaly
- Thunderstorm (temporal damage on one of proton 2023-6-29)
- Possible fear: E10 highway broadening


Use of data

- 2022-4-15 event (large SC-like ∆B without SW shock)
 - presented at EGU (April 2023)
- Kiruna ASC Auroral index
 - correlation to dB/dt is under way
- 2023-2-26 event (SC-driven shock aurora at 21 MLT)
 - correlate magnetic signature to different aurora
- Zeeman effect vs magnetic variation
 - effect changes with time (relation to magnetic variation?)



2022-4-15 event







filtered quick data at Kiruna UT dX dY d7 58.2 30.8 45.4 04:28:10 53136.0 04:30:13 0.7 22.2 53152.9 04:32:20 -342.2 36.4 63.2 53071.8 04:33:40 53163.6 -1.0101.5 88.6 53236.7 04:34:10 65.8 81.0


dX/dt = 50-60 nT/10s around the peak

backup magnetometer






Lycksele



global perspective (2022-4-15 event)

SuperMAG plot @ 04:33 UT



summary (2022-4-15 event)

- Large isolated ∆B skipe (~ 400 nT in Kiruna), localized @ dawn
- ∇× I_H kernel is moving ⇒ downward J_{//} moving toward night
- Riometer data is consistent with downward J_{II}
- At low latitudes, SC-like compression at wide longitude
- Unlikely SC, no substorm/storm is followed (rather ended)

- But no shock detection by any of L1 or MMS/Cluster.
- Stable northward $B_z > +10$ nT (L1 and MMS), but big dip at Cluster (~ to near zero) during the density enhancement.

⇒ Mysterious!




ASC auroral Index

 Just from ASC, Level 6 (brightening and breaking in the ASC), and Levels 4a+4b (possible precursor) are defined

Additional Level using |dB/dt|

- Level 8: |dB/dt| ≥ 5 nT/s
- Level 7: Level 6 & |dB/dt| > 2 nT/s
- Level 5: some ASC criteria + dB/dt > 0.8 nT
- Level 4c: some ASC criteria + dB/dt
 > 0.5 nT






Probability of reaching Level 6


from first time of reaching the lower Level over-night

Note: Preliminary result with insufficient statistics due to limited (November 2021 – April 2022).

cf. Level 8 (> 5 nT/s)

Too little statistics (only 6 cases of Level 8), and only 3 cases had precursor (Level 5, Level 6, and Level 7).







Correlation between |dB/dt| and ASC auroral index

Note: Preliminary result with insufficient statistics due to limited (November 2021 – April 2022).


UT	16-18 UT	18-22 UT	20-22 UT	22-24 UT
dB/dt vs %arc	0.46	0.27	0.54	0.41
dB/dt vs %strong	0.40	0.20	0.26	0.36
$ dB/dt $ vs $L_3 = aver(L^3)$	0.39	0.22	0.26	0.36

$$MLT \approx UT + 1.5 hr$$

- Correlation between the aurora activity (ASC auroral index) not very high, but still exists.
- Prediction of dB/dt > 5 nT/s (Level 8) is yet difficult, but Level 7 (Level 6 + |dB/dt| >2 nT/s) can be a possible candidate.
- We will examine more data in this direction (add 2022/2023 data and add 2023/2024 data).




shock aurora @ 21 MLT: 2023-2-26




DL and DP (primary and main) after SC

reversal of $\Delta H \rightarrow$ conversing $J_P \rightarrow$ upward field-aligned current








Kiruna


Sweden


Zeeman effect seen in mm wave (O₂ & O₃)




for further information, contact

uwe.raffalski@irf.se







End / questions?



Magnetometer quality


Scalar/vector difference may be due to time resolution difference, otherwise good agreement

Variometer

Proton (#2 & #3) & Variometer_F


Proton #2 – Variometer_F


Proton #3 - Proton #2





Absolute (baseline) measurement










Quality of 1-sec data





We calculate fluctuation of dB/dt

- a. 1-min average of |dB/dt|_1sec
- b. 3-hour standard deviation of |dB/dt|
- c. 1-month average of std(|dB/dt|)_3h

⇒ Good fit with other stations

The result is valid for both (DMI and DTU) magnetometers.






Mining effect?


Only if we make complicated analyses, we can see mining effect

- a. 1-min average of |dB/dt| 1sec
- b. take Z/H ratio of this 1-min average
- c. 3-hour standard deviation of this ratio
- d. 1-month average of std(ratio)_3h
- ⇒ We finally see Kiruna anomaly
- dZ/dt is relatively small compared to dH/dt
- · consistent with high conductivity.

The result is again valid for both (DMI and DTU) magnetometers.


std[|dZ/dt|/|dH/dt]_3h










Mining effect?





Mining effect?





$< std[(dZ/dt)/(dH/dt)]_3h >_month$

